



New Perspectives on the NEO Threat and Other Reasons to Consider Asteroid Deflection/Fragmentation

Mark Boslough
Sandia National Labs
Albuquerque, NM

Asteroid Deflection Research Symposium 2008
October 23-24, 2008

Red Storm simulation

2005 simulation was for concept demonstration: not intended to be realistic option

New perspectives on the impact threat

1. Some nuclear deflection strategies are better:
Multiple low-yield shallow bursts are best.
2. Low-altitude airbursts dominate residual threat:
Mitigation should be for ~hundred m asteroids.
3. Recent controversies in impact science:
Credibility of community requires solid research.
4. Total asteroid threat is diminishing:
Mitigation not justified for NEO protection alone.
5. Better reasons to consider asteroid deflection:
Geo-engineering and space resources.

1. Nuclear deflection

- Best way to couple energy and momentum is buried nuclear burst (direct consequence of scaling laws)
- For a given total yield, multiple bursts are better than single burst (direct consequence of scaling laws)
- Non-weaponizable impact-triggered fission device is possible in principle
- Impact-triggered fission device would make multiple low-yield shallow bursts feasible


Numerical simulations of momentum coupling

- Three prongs of analysis: Numerical simulations, analytic models, scaled experiments.
- 2D parameter studies, sensitivity analysis, and intuition (require workstation computing).
- 3D models quantify effects of heterogeneities, strength, fracture, and spallation that are not amenable to either scaled experiments or analytical models (require capability computing).



2D workstation-class problem

Example:

1 kiloton surface burst in an intermediate-strength 100-meter diameter dunite disk ($m=10^9$ kg) couples 3×10^8 kg·m/s, but spalls material off the back surface.




3D capability-class problem



Example:

For known 3D geometry, internal structure of Itokawa must be assumed. Over what range of parameters can momentum be coupled to Itokawa without breaking it?

Golevka deflection: 100 kt explosion, 5 m deep 1 billion computational cells, Red Storm




Time = 0.00000

2. Low-altitude airbursts

- The relative threat from low-altitude airbursts (LAAs) is increasing
- Our understanding of LAAs is improving
- The next destructive NEO will almost certainly be a LAA
- “100/100/100 event”: (~100 m, ~100 Mt, ~1/100 chance this century)
- 100/100/100 event will dominate threat after survey is complete
- Time for technology development is similar to threat reduction time
- There are other reasons to deflect/fragment small asteroids
- Technology should be focused small (~few hundred meter) asteroids


The nature of the impact threat is changing



There are two types of Low-Altitude Airburst

1. The explosion generates a fireball that descends rapidly but does not reach the ground. Most of the damage at ground level is mechanical, due to the blast wave. This occurs for explosions between about 1 and 10 Mt, and may occur on time scales of several hundred years. The only known example is 1908 Tunguska event.
2. The fireball is much larger and descends all the way to the surface. The damage is dominated by high-temperature thermal radiation. The threshold yield is about 10 Mt, depending on other parameters, and they recur on time scales of several thousand years. The best putative example is the Libyan Desert Glass event, 29 million years ago.

There are two types of Low-Altitude Airburst



Type 1: Tunguska

Type 2: Libyan Desert

Type 1 airburst simulation: 5 megaton

Type 2 airburst simulation: 15 megaton

Type 1 LAA: “Tunguska-Type”



Type 1 LAA: “Tunguska-Type”

Florenskiy: 1961 Expedition



Consequences of Type 1 airburst





Krinov, 1963




Fig. 485 — THE SIBERIAN FOREST DEVASTATED BY THE BLAST FROM THE METEORITE OF 30 JUNE 1908.




Modeling Type 1 airburst



5 Mt explosion at 12 km above surface, 35° entry angle



Tunguska treefall map (Longo et al, 2005)



Wind speed map (this study)



Type 2 LAA: “Libyan-Desert-Type”




Consequences of Type 2 airburst




Modeling Type 2 airburst



15 Mt explosion at 18 km above surface, 35° entry angle




Tunguska treefall map (Longo et al, 2005)



Wind speed map (this study)

100 Mt explosion, 90° entry angle

Temperature at 3.56 seconds




1 km

Temperature shading

100 Mt explosion, 90° entry angle

Wind Speed at 3.53 seconds



1 km

Wind speed shading

Animated version for National Geographic
documentary “Ancient Asteroid”

Tutankhamun's Fireball (Ancient Asteroid)

Meteor Crater impact was a low-altitude airburst

Less than 30% of the initial energy made it to the surface (Melosh & Collins, 2005)
9.0 Mt initial KE, 2.5 Mt formed crater, 6.5 Mt explosion in air




The physics of airbursts



Shoemaker-Levy 9 comet crash: Jupiter, 1994

Shoemaker-Levy 9 impact, July 1994




Visible From Earth


Behind Jupiter

This is a *robust* result

Shoemaker-Levy 9 impact, July 1994




Impact G



Hubble Space
Telescope Image

This is a *validated* result

View from Earth



Shoemaker-Levy 9 comet crash: Jupiter, 1994

Plumes and line explosions on Earth



**Annals of
the New York Academy
of Sciences
Volume 822**

Near- Earth Objects

**The
United Nations
International
Conference**

**Editor
John L. Remo**






FIGURE 16. Comparison of butterfly-shaped treefall pattern at Tunguska [17] to map of peak surface windspeeds calculated from the collision of a 12 megaton object entering the atmosphere at an oblique angle, 55° from the zenith.

3 MT impact source produces high velocity "ballistic fireball"



3 MT explosion source produces high velocity "buoyant fireball"




Pancake model: Earth's atmosphere protects us from low-altitude airbursts

Chyba et al. (1993), "The 1908 Tunguska explosion: atmospheric disruption of a stony asteroid" *Science*.

$$m \frac{dv}{dt} = -\frac{1}{2} C_D \rho A v^2$$



$$q \frac{dm}{dt} = -\frac{1}{2} C_H \rho A v^3$$

$$r \frac{d^2 r}{dt^2} = -\frac{1}{2} C_D \frac{\rho}{\rho_m} v^2$$



A stony asteroid deposits essentially all of its kinetic energy above 7 km. In this model the energy deposition curve is sharply peaked because of the mutually-reinforcing effects of atmospheric drag and deformation. Subsequent modeling has been based on point-source explosions and nuclear weapons effects.


Pancake model revisited: Earth's atmosphere is actually penetrated by hot vapor jet



The “point source explosion” model is a poor approximation.


Movies: Difference between explosion and impact

5 megaton point explosion at 5 km altitude: first 20 seconds



Movies: Difference between explosion and impact

5 megaton point explosion at 5 km altitude: first 20 seconds




Box dimensions: 8.4 x 15 km

Movies: Difference between explosion and impact


5 megaton impact airburst at 5 km altitude: first 20 seconds

Time = 10.02 seconds



Movies: Difference between explosion and impact

5 megaton impact airburst at 5 km altitude: first 20 seconds




Box dimensions: 8.4 x 15 km

Impact-Induced Vortex Rings

5 megaton impact airburst at 5 km altitude
generates stack of supersonic white-hot “mega-tornado” rings

← 4 km →



4 km

Red: inward rotation
Blue: outward rotation

Sustained supersonic flow
melts and ablates surface

Impact-Induced Vortex Rings


5 megaton impact airburst at 5 km altitude





Box dimensions: 3 x 4 km

Bologna, Italy 1996

Tunguska96



Silica96



Animated version for National Geographic documentary
“Ancient Asteroid”

This video was shown as justification for Younger-Dryas extinction impact hypothesis at 2007 Spring AGU, Acapulco:

<http://www.youtube.com/watch?v=I2ld-lohrPw&feature=PlayList&p=741568C2D58A9793&index=1>

This idea is based on simulations of Boslough and Crawford (1997) but is not physically realistic as shown in this animation. There is no viable mechanism to generate a cluster of fragments with this spacing.

Tutankhamun's Fireball (Ancient Asteroid)

3. Recent impact controversies

Two new papers



Gasperini et al. (2007) *“A possible impact crater for the 1908 Tunguska Event”*

- Viable hypothesis
- Only evidence is circumstantial
- Have not found any theoretical support

Firestone et al. (2007), *“Evidence for an extraterrestrial impact 12,900 years ago that contributed to the megafaunal extinctions and the Younger Dryas cooling”*

- Not a viable hypothesis
- Evidence is very weak
- Completely inconsistent with physics of impacts and airbursts

The Lake Cheko Controversy



"A possible impact crater for the 1908 Tunguska Event" Gasperini et al., 2007

The Lake Cheko Controversy

Lake Cheko photography:
Discovery Channel


Impact crater or just a lake?




First post-expedition conference: Vanavara

First look: Submersible camera videos!

The Lake Cheko Controversy



“Anfinogenov spindles” (Anfinogenov, 1966)



Multiple explosions (Boslough, 2008)

Have not been able to find a physically-reasonable model that creates airburst consistent with Tunguska treefall pattern and crater consistent with Lake Cheko. Airburst modeling does not disprove the hypothesis, but does not support it either.

The Holocene Impact Rate Controversy



June 2008 Atlantic Monthly

“The odds that a potentially devastating space rock will hit Earth this century may be as high as one in 10. So why isn’t NASA trying harder to prevent catastrophe?”

By **Gregg Easterbrook**


A team of researchers led by **Richard Firestone**, of the Lawrence Berkeley National Laboratory, in California, recently announced the discovery of evidence that one or two huge space rocks, each perhaps several kilometers across, exploded high above Canada 12,900 years ago.

If, as **Boslough** thinks, most asteroids and comets explode before reaching the ground, then this is another reason to fear that the conventional thinking seriously underestimates the frequency of space-rock strikes—the small number of craters may be lulling us into complacency.

Given the scientific findings, shouldn’t space rocks be one of NASA’s priorities? You’d think so, but **Dallas Abbott** says NASA has shown no interest in her group’s work: “The NASA people don’t want to believe me. They won’t even listen.”

The Atlantic article was very misleading

The Megafauna Extinction Controversy



Firestone et al., 2007
(poster, AGU Fall Meeting)




R.B. Firestone et al. (2007),
"Evidence for an extraterrestrial impact 12,900 years ago that contributed to the megafaunal extinctions and the Younger Dryas cooling"



Bear skull, Krasnoyarsk

Not consistent with physics of airbursts

next figure abnormal splash fire wood (Weiss, 1987) common
4000-10000 years ago.
Heroes of tragedy




Russian poster at
Tunguska 2008
conference, Moscow

4. The asteroid threat in perspective



The asteroid threat in perspective


Alan Harris,
 “What Spaceguard Did”
Nature 453, 2008



Cause	Expected deaths per year*
Type 1 airburst	10 (<1 per year in US)
All asteroids	80 (~4 per year in US)
Climate change	150,000 – 300,000
Air Pollution	600,000
Malaria	1,000,000
Traffic accidents	1,200,000
HIV/AIDS	2,100,000
Tobacco	5,000,000

*Asteroid estimates from Harris (2008).
 All other estimates from the World Health Organization.


5. Better reasons for NEO deflection



The rate of climate change is exceeding the most extreme expectations:
Huge uncertainty in climate projections is analogous to uncertainty in NEO orbits


Climate change threat is ~ 10,000 times greater

...when probability distribution function for climate sensitivity is treated the same way as probability distribution function for NEO impact



$$\text{Threat} = \text{Probability} \times \text{Consequences}$$

NEO capture for space resources or geo-engineering





Low-energy pathways allow capture of object into Earth orbit (Marsden and Ross, 2005)

Global climate change mitigation: More urgent than impact mitigation by $\sim 10^4$

- Captured asteroids can be fragmented and used to create engineered orbital debris ring, or placed at L1 for shadow.
- Changes in incoming solar radiation and atmospheric opacity modify Earth's climate.
- Rapid changes in Earth's climate are observable in geologic record.



Climate simulations



Difference in mean temperature
(low vs. high obliquity)

Conclusions

1. Multiple low-yield shallow bursts are most effective nuclear deflection method.
2. Primary mitigation goal should be for Type 2 airburst prevention.
3. Credibility of community requires skepticism and solid research.
4. Deflection not justified for NEO protection alone.
5. Geo-engineering and space resources are vastly better reasons to develop deflection technology.

Questions?

